



UNIVERSITI PUTRA MALAYSIA

**OPTIMISATION OF ENZYME AIDED PEELING OF MUSK LIME
(*CITRUS MITIS* B.) AND DEVELOPMENT OF ITS CANDIED PEEL**

HAZNIZA DATO' HJ. ADNAN

FSMB 2003 30

**OPTIMISATION OF ENZYME AIDED PEELING OF MUSK LIME
(*CITRUS MITIS* B.) AND DEVELOPMENT OF ITS CANDIED PEEL**

HAZNIZA DATO' HJ. ADNAN

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2003



**OPTIMISATION OF ENZYME AIDED PEELING OF MUSK LIME
(*CITRUS MITIS* B.) AND DEVELOPMENT OF ITS CANDIED PEEL**

By

HAZNIZA DATO' HJ. ADNAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

October 2003



To my beloved families and friends....

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Master of Science

**OPTIMISATION OF ENZYME-AIDED PEELING OF MUSK LIME
(*CITRUS MITIS* B.) AND DEVELOPMENT OF ITS CANDIED PEEL**

By

HAZNIZA DATO' HJ. ADNAN

October 2003

Chairman : Associate Professor Azizah Osman, Ph.D.

Faculty : Food Science and Biotechnology

Studies on the optimisation of enzyme peeling of citrus fruits were carried out on musk lime (*Citrus mitis* B.). The time taken to peel musk lime by enzyme (Peelzym II) was minimised with increasing enzyme concentration and application of vacuum pressure. The optimum enzyme concentration and optimum vacuum pressure for peeling musk lime peel was 1.0% (v/w) and 700 mm Hg, respectively. The effect of vacuum pressure in physico-chemical changes of peeled fruits of musk lime (*Citrus mitis* B.) during enzymatic peeling was studied. The pH, total soluble solids, citric acid, ascorbic acid, moisture, total pectin, tannin and sugar contents in the puree of peeled musk lime were not significantly affected by the vacuum pressure, except, total pectin content and 'b' value of puree colour at very high vacuum pressure (700 mm Hg). The naringin content has also been analysed in the puree, seeds and peel (discarded) of peeled musk lime. The

lime. The naringin content in the puree and seeds of peeled musk lime were not significantly affected by the vacuum pressure, however, naringin content in the peel was significantly affected at very high vacuum pressure. The development of candied musk lime peel was carried out using Response Surface Methodology (RSM) and ECHIP[®] software. In this study, the RSM method was found to be effective in making a new product when the developed product (candied musk lime peel) had matched the sensory profile of a target product (commercial candied peel). This method was then confirmed by a verification process between the experimental value from the developed product (candied musk lime peel) with the target value obtained from the overlapping contour maps of ingredients in the making of optimum candied musk lime peel. Results from sensory evaluation also showed that most of the panelists liked the taste of the product compared to other sensory attributes tested. In comparison, the developed candied musk lime peel has a similar profile in terms of appearance, odour and taste of sourness and bitterness to that of commercial candied peel. Therefore, the candied musk lime peel developed in this study has the potential to be commercialised.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGOPTIMUMAN PROSES PENGUPASAN BUAH LIMAU KASTURI
(*CITRUS MITIS* B.) TERBANTU ENZIM DAN PENGEMBANGAN HALWA
DARI KULITNYA**

Oleh

HAZNIZA DATO' HJ. ADNAN

Oktober 2003

Pengerusi : Profesor Madya Azizah Osman, Ph.D.

Fakulti : Sains Makanan dan Bioteknologi

Kajian ke atas pengoptimuman proses pengupasan buah sitrus terbantu enzim telah dijalankan ke atas buah limau kasturi (*Citrus mitis* B.). Masa yang diambil untuk buah limau kasturi dikupas oleh enzim (Peelzym II) telah diminimakan dengan meningkatkan kepekatan enzim dan tekanan vakum yang digunakan. Kepekatan optimum enzim dan tekanan optimum vakum untuk pengupasan buah limau kasturi ialah 1.0% (isipadu/berat) dan 700 mm Hg, masing-masing. Kesan tekanan vakum dalam perubahan fisiko-kimia buah limau kasturi (*Citrus mitis* B.) yang telah dikupas semasa pengupasan terbantu enzim telah dikaji. Nilai pH, jumlah pepejal larut, kandungan asid sitrik, asid askorbik, kelembapan, jumlah kandungan pektin, tanin dan gula di dalam puri limau kasturi yang telah dikupas adalah tidak signifikan terhadap kesan tekanan vakum, kecuali, jumlah kandungan pektin dan nilai 'b' bagi warna puri pada kesan tekanan vakum yang tinggi (700 mm Hg). Kandungan naringin juga telah dianalisa di dalam

kecuali, jumlah kandungan pektin dan nilai 'b' bagi warna puri pada kesan tekanan vakum yang tinggi (700 mm Hg). Kandungan naringin juga telah dianalisa di dalam puri, biji dan kulit (hampas) limau kasturi yang telah dikupas. Kandungan naringin di dalam puri dan biji bagi limau kasturi yang telah dikupas adalah tidak signifikan terhadap kesan tekanan vakum, tetapi, kandungan naringin di dalam kulit adalah signifikan pada kesan tekanan vakum yang tinggi. Pengembangan hasil bagi halwa kulit limau kasturi telah dijalankan dengan menggunakan kaedah 'Response Surface Methodology' dan software ECHIP[®]. Di dalam kajian ini, kaedah RSM didapati sangat berkesan dalam menghasilkan produk baru apabila produk yang dihasilkan (halwa kulit limau kasturi) sepadan dengan profil ujirasa produk sasaran (halwa kulit limau komersial). Kaedah ini kemudiannya telah diperakui melalui proses pengesahan diantara nilai ujikaji yang didapati daripada produk yang dihasilkan (halwa kulit limau kasturi) dengan nilai sasaran yang diperolehi daripada penindihan peta kontur ramuan-ramuan di dalam penghasilan halwa kulit limau kasturi yang optimum. Keputusan ujirasa juga menunjukkan kebanyakan panelis menyukai rasa produk berbanding lain-lain ciri ujirasa yang diuji. Secara bandingan, halwa kulit limau kasturi yang dihasilkan mempunyai persamaan profil dari segi rupabentuk, bau dan rasa dengan halwa kulit limau komersil. Maka, halwa kulit limau kasturi yang telah dihasilkan di dalam kajian ini didapati berpotensi untuk di komersilkan.

ACKNOWLEDGEMENTS

Alhamdulillah, first of all I would like to express my utmost thanks and gratitude to the Almighty Allah, the Sustainer, the Most Gracious and Most Merciful, without whose will no one can achieve anything, and my salawat and salam to His righteous messenger, prophet Muhammad S.A. W.

I would like to take this opportunity to express my appreciation and gratitude to the chairman of my Supervisory Committee, Assoc. Prof. Dr. Azizah Osman for her invaluable suggestion, guidance, discussion and patience throughout the project. I am also grateful to the other member of my supervisory committee, Prof. Dr. Hasanah Mohd. Ghazali and Assoc. Prof. Dr. Russly Abdul Rahman for their constructive comments towards the project and preparation of this thesis. My sincere gratitude is also extended to the financial support provided by the IRPA fund for this research, which was awarded to Assoc. Prof. Dr. Azizah Osman. I am also indebted to all the staffs of the Department of Food Science, Biotechnology and Food Technology for their generous cooperation. Acknowledgements are also due to all my labmates, friends and also laboratory assistants who had given me the moral encouragement and support to complete my graduate study.

Finally, I also wish to express my deepest appreciation to my beloved husband and daughters, my parents and my parent in-laws, who have given me encouragement and morally support in anyway during the many years of my seemingly never ending pursue for knowledge.

I certify that an Examination Committee met on (date of viva) to conduct the final examination of Hazniza Dato' Hj. Adnan on her Master of Science thesis entitled "Optimisation of Enzyme-Aided Peeling of Musk Lime (*Citrus mitis* B.) and Development of Its Candied Peel" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

AZIZAH ABDUL HAMID, Ph.D.


Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

AZIZAH OSMAN. Ph.D

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

RUSSLY ABDUL RAHMAN. Ph.D

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)



.....
GULAM RUSUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11 FEB 2004

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows:

AZIZAH OSMAN. Ph.D

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

HASANAH MOHD. GHAZALI. Ph.D

Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

RUSSLY ABDUL RAHMAN. Ph.D

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : **25 FEB 2004**

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citation, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or other institutions.



.....
HAZNIZA DATO' HJ. ADNAN

Date: 20 JAN. 2004

TABLE OF CONTENTS

	Page
DEDICATION.....	ii
ABSTRACT.....	iii
ABSTRAK.....	v
ACKNOWLEDGEMENTS.....	vii
APPROVAL	viii
DECLARATION FORM.....	x
LIST OF TABLES.....	xv
LIST OF FIGURES	xvii
LIST OF PLATES	xviii
 CHAPTER	
1 INTRODUCTION	1
 2 LITERATURE REVIEW	5
Citrus Fruits.....	5
Physico-chemical Characteristics of Citrus Fruits.....	9
Juice Colour.....	9
Acidity.....	10
Ascorbic Acid.....	11
Total Soluble Solids (TSS).....	11
Moisture Content.....	12
Pectic Substances.....	13
Sugar Content.....	13
Tannin Content.....	14
Naringin Content.....	15
Fruit Processing.....	18
Peeling Technology.....	19
Conventional and Lye Peeling.....	19
Fruit Pretreatment.....	20
Pectic Enzymes.....	21
Peelzym.....	22
Enzyme Aided Peeling.....	23
Vacuum Infusion Process.....	24
Quality of Peeled Fruit.....	25
By-products Development	25
Discarded Peel	26
Candied Peel.....	27
Optimisation Process for By-product development.....	28
Sensory Evaluation.....	29

3 OPTIMISATION OF ENZYMATIC PEELING CONDITIONS AND PHYSICO-CHEMICAL CHARACTERISTION OF PEELED MUSK LIME (*CITRUS MITIS* B.)

Introduction	32
Materials and Methods	34
Source of Fruits.....	34
Preparation of Enzyme Solution.....	34
Preparation of Fruits.....	35
Effect of Enzyme Infusion Methods on Peeling.....	35
Determination of Optimum Enzyme Concentration.....	38
Determination of Optimum Vacuum Pressure.....	38
Preparation of Samples.....	39
Physico-chemical Analysis of Treated Musk Lime.....	41
Puree Colour.....	41
Titratable Acidity (TA).....	42
pH.....	42
Ascorbic Acid Content.....	43
Total Soluble Solids (TSS) Content.....	44
Moisture Content.....	44
Sugar Content.....	45
Total Pectin Content.....	46
Tannin Content.....	47
Naringin Content.....	48
Statistical Analysis.....	48
Results and Discussion	50
Optimisation of Enzyme Aided Peeling.....	50
Enzyme Infusion Method.....	50
Optimum Enzyme Concentration.....	51
Optimum Vacuum Pressure.....	54
Changes in the Physico-chemical Characteristics of Peeled Musk Lime.....	56
Physico-chemical Characteristics.....	59
Puree Colour.....	59
pH, Titratable Acidity and Ascorbic Acid Content.....	62
Total Soluble Solids (TSS) and Moisture Content.....	64
Total Pectin and Tannin Contents.....	66
Sugar and Total Sugar Contents.....	68
Naringin Content.....	70
Conclusion	72

4 DEVELOPMENT OF CANDIED MUSK LIME PEEL

Introduction	74
Materials and Methods	75
Source of Peel.....	75
Raw Materials.....	76
Processing of Candied Musk Lime Peel.....	76
Preparation of Candied Musk Lime Peel.....	76
Experimental Design.....	78
Sensory Evaluation of Candied Musk Lime Peel.....	79
Verification of Optimum Candied Musk Lime Peel.....	82
Physico-chemical Analysis of Optimum Candied Musk Lime Peel.....	83
Peel Firmness.....	83
Chemical Analysis.....	83
Statistical Analysis.....	84
Results and Discussion	85
Sets of Formulation by Applied RSM.....	85
Training of Selected Sensory Panelists.....	87
Regression Analysis on Sensory Attributes.....	88
Development of Candied Musk Lime Peel.....	90
Optimisation of Candied Musk Lime Ingredients.....	90
Verification of Optimum Formulation (Target Value)....	92
Descriptive Panel Analysis.....	95
Acceptance Test.....	98
Conclusion	105

5 CONCLUSIONS AND RECOMMENDATIONS

Conclusions	106
Recommendations	107

REFERENCES.....	108
------------------------	------------

APPENDICES

A (Additional Plates).....	119
B (Schematic Diagram of Musk Lime Fruit).....	128
C (Additional Tables).....	130
D (Questionnaire-Sensory Sheet).....	133
1 (Screening Part I).....	134
2 (Screening Part II).....	135
3 (Training Part I).....	137
4 (Training Part II).....	138
5 (Sensory Terminology).....	139
6 (Selection of Commercial Candied Peel as a Reference Sample).....	140

7 (Hedonic Test and Qualitative and Descriptive Analysis I).....	141
8 (Hedonic Test and Qualitative and Descriptive Analysis II).....	144
9 (Hedonic Test and Qualitative and Descriptive Analysis III).....	147
10 (Hedonic Test - Verification).....	150
11 (Ranking for Preference).....	151
BIODATA OF AUTHOR.....	152

LIST OF TABLES

Table		Page
1	Physico-chemical Composition of Musk Lime Fruits and Related Findings.....	57
2	Effect of Different Vacuum Aided Enzyme Infusion on the Changes In Colour (Hunter 'L', 'a' and 'b') of Peeled Musk Lime.....	60
3	Effect of Different Vacuum Aided Enzyme Infusion on the Changes in pH, Titratable Acidity (TA) and Ascorbic Acid (AA) Content of Peeled Musk Lime.....	63
4	Effect of Different Vacuum Aided Enzyme Infusion on the Changes in Total Soluble Solids (TSS) and Moisture Content of Peeled Musk Lime.....	65
5	Effect of Different Vacuum Aided Enzyme Infusion on the Changes in Total Pectin Content and Tannin Content of Peeled Musk Lime	67
6	Effect of Different Vacuum Aided Enzyme Infusion on the Changes in Fructose, Glucose, Sucrose and Total Sugar Content of Peeled Musk Lime	69
7	Effect of Different Vacuum Aided Enzyme Infusion on the Changes in Naringin Content (Peel, Seeds and Puree) of Peeled Musk Lime....	71
8	A list of Descriptor and Definition Used in QDA Test.....	81
9	Formulation Designed by RSM to Verify the Amount of Ingredients (Sugar syrup, Sorbitol and Citric Acid) Used in the Development of Candied Musk Lime Peel.....	86
10	Significance of the Factors and the Interactions Between the Ingredients Coefficient and Sensory Variables.....	89
11	Hedonic Profiles of Each Sensory Attribute for Candied Musk Lime Peel (CMLP).....	94
12	QDA Profiles of Candied Musk Lime (CMLP) and Commercial Candied Peel (CCP).....	96

13	Hedonic Results for Acceptance Profiles of Candied Musk Lime Peel (CMLP) and Commercial Candied Peel (CCP).....	100
14	Analytical Data for Candied Musk Lime Peel (CMLP) and Commercial Candied Peel (CCP).....	104
15	Enzyme Activities of Peelzym I,II,III and IV as Reported in the Product Sheet.....	131
16	Preparation of Enzyme (Peelzym II) Concentration, (v/w, %) where Volume of Enzyme Used Based on 900 g of Fruits.....	131
17	Determination of the Best Technique in Enzyme Infusion Method Between Method I (Immersion in Enzyme Solution) and Method 2 (Suck in Enzyme Solution).....	132

LIST OF FIGURES

Figure		Page
1	Cross Section of a Citrus Fruit.....	6
2	Structure of Naringin (Naringenin 7 β -neohesperidoside).....	17
3	Preparation of Peeled Musk Lime Samples.....	40
4	Extraction and Preparation of Naringin Sample.....	49
5	Peeling Efficiency of Different Enzyme Concentrations.....	52
6	Peeling Efficiency of Different Vacuum Pressures.....	55
7	Production of Candied Musk Lime Peel.....	77
8	Effect of Ingredient Levels on Correx Reading.....	91
9	The Overlaid Contour Maps of Five Dependent Sensory Attributes of Candied Musk Lime Peel.....	93
10	Cobweb Plot of Sensory Profiles of Candied Musk Lime Peel and Commercial Candied Peel for Appearance (Colour, Moist-Dry), Odour (Sweet, Sour), Taste (Sweetness, Sourness, Bitterness) and Texture (Hardness, Tear Ability) Attributes	97
11	Acceptance Profiles on Five Sensory Attributes for Candied Musk Lime Peel.....	99
12	Acceptance Profiles on Taste of Candied Musk Lime Peel.....	101
13	Acceptance Profiles on Sensory Attributes of Candied Musk Lime Peel (a) and Commercial Candied Peel (b).....	103
14	Schematic Diagram of Flavedo-Scouring of Musk Lime at Five Different Parts of Its Surface.....	129

LIST OF PLATES

Plate		Page
1	Diagrams for Method 1.....	36
2	Diagrams for Method 2.....	36
3	A Tree of Musk Lime (<i>Citrus mitis</i> B.) Fruit.....	120
4	A Thornless Branch of Musk Lime (<i>Citrus mitis</i> B.) Tree.....	120
5	A Matured Musk Lime (<i>Citrus mitis</i> B.) Fruits	121
6	A Batch of Musk Lime Fruits (100 g Equivalent to 8-10 Fruits).....	121
7	A Batch of Scoured Musk Lime Fruits Using an Abrasive Paper.....	122
8	Enzymatic-Peeling Process in Shaking Incubator.....	122
9	A Peeled Musk Lime Fruit (After Enzyme and Vacuum Pressure Treatment).....	123
10	A Peeled Musk Lime Fruit (Hand Peeling).....	123
11	Discarded Peel of Musk Lime Peel – A By-Product of Enzymatic Peeling.....	124
12	Musk Lime Soaked in Brine Solution During Pickling Process.....	124
13	Musk Lime Soaked in Syrup Solution During Candying Process.....	125
14	Musk Lime Peels Arranged in Wooden Tray After Candying Process..	125
15	Cabinet Dryer for Drying the Musk Lime Peels.....	126
16	Musk Lime Peels After Drying Process.....	126
17	Musk Lime Peels After Coated With Sugar.....	127
18	Commercial Candied Peel as a Reference Sample in Sensory Evaluation.....	127

CHAPTER 1

INTRODUCTION

Citrus which originated from Southeast Asia, is one of the most important fruit crops in Asia (Ko, 1996). However, yields per unit area in Asia are low compared to most Western countries. Citrus has been chosen as one of the 16 fruit types for commercial production under the Malaysian National Agricultural Policy (Ko, 1996). Presently, there are about 1600 species of citrus in the world amongst which only a few of the better-known species are widely cultivated in Malaysia. The main citrus cultivated are oranges, pomelos and limes (Faridah and Azizah, 1982). In Malaysia, domestic production area of citrus fruit varies considerably between each fruit type. Lime is the second most important citrus after pomelo, followed by sweet orange and mandarin (limau langkat).

To date, reports on the quality of commercial musk lime juice are not available. This may be due to the once less popular status of commercially musk lime juice compared to orange juice. At present, there are many problems confronting the production of musk lime juice. In the industry, juice extraction using screw press is considerably impractical for the extraction of musk lime juice (Swisher and Swisher, 1977). In the process, the whole fruit is crushed (by pressure) in a screw press, then pass through a centrifugal machine to separate the juice from the mixture of peel and pulp (Kale and Adsule, 1995). This excessive extraction pressure could split the seeds and break the peel, which generally contain the bitter principle (naringin) and could also

release excessive peel oil into the juice (Jainudin and Mazuin, 1988). Hence, the existence of this bitter taste in the juice is the result of the processing technique. The pronounced bitter taste was relatively less preferable to most of the consumers. The physiological limitation of musk lime fruit such as its small size and very thin peel compared to other citrus fruits, makes the mechanical juice extraction unsuitable for the extraction of musk lime juice. An understanding of these problems is, therefore, of the utmost importance in the promotion of the industrial value of the musk lime juice.

Conventional methods for fruit peeling such as hand peeling, mechanical, chemical, acid, heat and flame usually produce 30-40% of edible waste (Baker and Grohmann, 1995). This waste, containing live cells, are sensitive to the environment and continue to change after processing due to chances of contamination with microorganisms (Setty et al., 1993). However, this wastage can be reduced by recycling (enzyme solution) processes if a proper processing technique is carried out. Therefore, in order to reduce the undesirable effects and the peeling losses due to the conventional peeling methods, enzymatic peeling was developed.

Enzymatic peeling has been defined as the application of exogenous enzymes specifically to alter the characteristics of intact tissues (McArdle and Culver, 1994). In the enzymatic peeling process, enzyme is made to infuse into intact plant tissues such as flavedo and albedo and effectively alter the functional properties of these tissues such as softening the fruit peel and making it feasible to remove (Bruemmer, 1981). The first macerating enzymes were tried on fruit mashes in the 60s mainly on blackcurrant due to the difficulties in extracting its juice (Janser, 1997). Today the maceration of fruit with

enzyme preparations is widespread and the advantages have been demonstrated for several fruits, conditions and extraction systems (Janser, 1997). The application of this technology was first developed for the production of fresh peeled fruit, fruit salads and segments. In 1981 a USDA patent was granted for the so-called vacuum infusion process with enzymes. Since then, a lot of development has been done (Janser, 1996).

In enzyme peeling process, a dilute solution of enzyme will be infused into intact citrus fruit tissues where the enzyme will penetrate between the citrus fruit segments loosening adhesion of the capillary membrane and allow the fruits peel to be easily removed (Janser, 1996). The enzymatic technology results in segments with improved freshness, better texture and appearance compared to traditional segments, which are peeled using a caustic soda process (Janser, 1997). Enzyme peeling of citrus benefits the processors by yielding 100% of the edible portion of fruit as salable product compared to the conventional peeling methods (Baker and Grohmann, 1995).

There are many methods that have been reported on ways to infuse enzyme into the fruit. One of the methods is the vacuum infusion where fruits are subjected to vacuum for a fixed time in an enzyme preparation for the infusion of the solution into the albedo layer and incubated in the enzyme preparation after breaking the vacuum until easy removal of the peel was obtained (Soffer and Mannheim, 1994). The advantages of fruit peeling by vacuum-oriented enzyme infusion are obvious. The quality of the obtained peeled fruit was better such as it had a more attractive appearance such as clean flesh, without white adhering albedo and more intensive colour and taste (Janser, 1996). This process also effectively alters the functional properties of the tissues

such as softening the fruit peel, firming the fruit flesh and enhancement of desired flavours (Elliott and Julia, 1993).

Fruit processing industries usually have to cope with 40-60% of the incoming harvested fruit exiting the juicers as waste (Baker and Grohmann, 1995). In the enzymatic peeling process, the loosened peel is considered as a waste. In attempting to produce by-products from the waste, the loosened peel has a high potential as a starting material for a development of value-added products. The loosen peel can be converted into more useful by-products through various processing technologies. The loosen peel can be candied, brined or pickled. The other citrus by-products that can be produced are bioflavonoids, flavourings and a gelling agent (Jainuddin and Mazuin, 1988).

Enzyme application for peeling fruits is new in Malaysia. Enzymatic peeling was introduced in Malaysia in late 1990's due to processing and market demand on local citrus juice and minimally processed citrus fruits. Since then, many research have been conducted to solve the processing problems, thus fulfill the market demand (Liu et al., 1999; Aziz et al., 1999).

Therefore, the objectives of this study were:

1. To determine the optimum conditions of enzymatic peeling of musk limes.
2. To determine the physico-chemical characteristics of enzymatic peeled musk limes.
3. To develop candied musk lime peel.

CHAPTER 2

LITERATURE REVIEW

Citrus Fruit

Citrus fruits are classified under the family Rutaceae and genus *citrus*, but botanically they are classified as berries. They are superior fruits with all of their tissues derived from the ovary, in contrast to apples in which some of the tissues are derived from the enlargement of the calyx and receptacle (Albrigo and Carter, 1977). The citrus fruit consists of an outer peel which serves largely as a cover to the inner portion. The outer portion of the peel is called the flavedo and the inner white spongy layer of parenchymatous cells is called the albedo. The flavedo contains numerous oil vesicles and chromatophores. The thickness of the albedo varies from species to species. The thickness of the albedo for the musk lime and mandarin orange is relatively thin measuring about 1.5 to 2.0 mm, while in the mexican lime is 2.5 to 3.5 mm (Jainudin and Mazuin, 1988). The edible portion of the fruit, the pulp is found in segments situated below the albedo layer. The structure (cross section) of citrus fruit is shown as in Figure 1.